


April 2001

Dairy Lines



Volume 7, Number 4

Editors

John F. Smith
*Extension Specialist
Dairy Science*

Dan Waldner
*Extension Specialist
Dairy Science*

Mike Brouk
*Extension Specialist
Dairy Science*

Contributors

Karen Schmidt
Professor, Dairy Products

John Shirley
Professor, Dairy Science

Jeff Stevenson
Professor, Dairy Science

Dave Sukup
Manager, Heart of America DHI



Printing sponsored by

MONSANTO
Food • Health • Hope™



DAIRY RESEARCH & EXTENSION NEWS

<http://www.oznet.ksu.edu/ansi/nletter/dairylin.htm>

Managing Milk Composition:

Feed additives can enhance health, growth and milk yields

This article is part of a fact sheet series produced by extension dairy specialists from Oklahoma, Texas and New Mexico.

Feed additives and management tools such as bST (bovine somatotropin) play an essential role in enhancing production and yield of milk and milk components. The need for a particular feed additive and its effectiveness depend upon a variety of factors. Producers are encouraged to critically evaluate the cost-to-benefit ratio of each feed additive in their management systems.

Feeding strategies that optimize rumen function result in maximum milk production and milk component percentages and yield. Additionally, producers who use records such as those provided by DHIA (Dairy Herd Improvement Association) can critically evaluate their nutrition and feeding-management programs.

Feed Additives

Supplemental Fat

Adding supplemental fat to rations for high-producing dairy cows has become a common practice. It is necessary to follow certain guidelines when feeding fat to cattle to avoid a depression of 0.1 to 0.2 units in the milk protein level. When used properly, added fat usually maintains or slightly increases milk fat percent, makes relatively little change in milk protein percent, and increases milk production. The net result is increased production of milk protein and nonfat solids. Too much fat in the ration can interfere with fiber digestion, reducing milk fat levels.

Limit total fat to 7.5 percent of the ration dry matter. A good rule of thumb is to

provide the same amount of fat in the ration as pounds of milk fat produced.

For example: 100 pounds of milk per day x 4.0 percent milk fat = 4 pounds of milk fat or 4 pounds total fat in the ration. Provide one-third of fat in the ration from normal ration ingredients, one-third from oilseeds or natural fats and one-third from rumen inert fat. Recommended guidelines for feeding fat are provided in Table 1.

Table 1. Fat Feeding Guidelines

Recommended Source	Maximum Inclusion
Basal diet	3%
Natural fats	2% - 4%
Whole oilseeds	1 lb.
Tallow	1 lb.
Protected fats	2% (1 lb.)
Total	6 - 7%

Note: When feeding supplemental fats, calcium and magnesium should be provided at 1.0 and 0.35 percent of the ration dry matter, respectively, because these fats can bind with calcium and magnesium and reduce their availability.

Sulfur

Sulfur is necessary for the synthesis of essential amino acids by rumen microbes. Sulfur supplementation is important in rations that contain high levels of nonprotein nitrogen (i.e., urea). Low sulfur intake can induce protein deficiency. The likelihood of this problem occurring increases with rations containing corn silage or poor quality grass silage. The recommended level of sulfur is 0.22 to 0.25 percent of the total ration dry matter.

Buffers

Buffers added to the diet help reduce the acid load placed on the rumen when high

continued on page 2

Continued from page 1

levels of grain are fed or when hay and grain are fed separately. Sodium bicarbonate, magnesium oxide, or a combination are the primary buffers recommended for feeding lactating dairy cows. Supplements of sodium bicarbonate should be 0.6 to 0.8 percent of the total diet dry matter or 1.2 to 1.6 percent of the concentrate mixture. Magnesium oxide should be added as 0.2 to 0.4 percent of the total diet dry matter or 0.4 to 0.6 percent of the concentrate mixture.

When feeding a combination of sodium bicarbonate and magnesium oxide, two to three parts sodium bicarbonate should be mixed with one part magnesium oxide and fed as a supplement at 0.8 to 1.2 percent of the total diet dry matter or 1.6 to 2.2 percent of the concentrate mix. Force-feeding larger amounts of these buffers may depress feed intake. Providing additional sodium bicarbonate free choice, beyond that which is already provided in the base ration, may prove beneficial in some herds when feeding for specific milk component changes. Estimated cost is 6 cents per head per day. The benefit-to-cost ratio is 4-to-1.

Rumen-protected amino acids

Responses to feeding individual amino acids to dairy cattle have not been consistent. Response differences probably occur based on the quantity and proportion of amino acids in the microbial and dietary protein digested and absorbed from the small intestine. Responses are often greater when mixtures of amino acids, rather than individual amino acids, are taken in beyond the rumen. Combinations of rumen-protected methionine and lysine have been shown to increase milk protein yield and concentration in diets low in rumen-degradable protein. Further, supplementing diets that contain added fat with rumen-protected methionine and lysine alleviates the milk protein depression effect of feeding added fat.

Yeast culture/fermentation products

Yeast culture and their fermentation products stabilize the rumen environment and improve fiber digestion. They maintain or increase dry matter intake and milk fat percent. Most benefits are seen in high producing cows or cows in early lactation. Feeding rate is 10 to 120 grams depending on yeast culture concentration. The cost is approximately 4 to 6 cents per cow per day. The estimated benefit-to-cost ratio is 4-to-1.

Niacin

Niacin, a water-soluble vitamin, was assumed to be produced in sufficient quantities by rumen microbes to meet the needs of the host animal. However, bacterial synthesis of niacin may not be adequate for high-producing cows. Milk yield and composition responses to niacin feeding are variable, at best. However, in some situations, niacin fed at 6 to 12 grams per day improves the milk protein depression caused by feeding high levels of fat. The estimated benefit-to-cost ratio is 6-to-1 for the 6-gram feeding level. The approximate cost is 1 cent per gram.

Bovine Somatotropin (bST)

The gross composition of milk (fat, protein and lactose) is not affected by treatment with bST. The factors that affect fat and protein content of milk of non-bST-treated cows have the same effects on milk composition of bST-treated cows. For example, certain breeds have a higher milk fat content, and an increase in milk fat typically occurs in late lactation for all breeds. Treat-

ment with bST does not alter these relationships. Likewise, the increase in milk fat content that occurs when the cow is using more energy than it is consuming and the decrease in milk protein content that occurs when the cow has an inadequate protein intake are also observed in bST-treated cows.

Milk from bST-treated cows also does not differ in vitamin content or in concentrations of nutritionally important mineral elements. In addition, proportions of total milk proteins represented by whey proteins and the different casein fractions are not changed substantially. Thus, the manufacturing characteristics are not altered by the use of bST to enhance milk yield in lactating cows.

Evaluating Cost Effectiveness

Consider the following factors in determining if a feed additive should be used:

- anticipated response
- economic return
- available research
- field response.

Anticipated response refers to performance changes such as increased milk yield, increased milk components, improved dry matter intake, improved growth, improved health, and/or minimized weight loss that could be expected when a feed additive is included.

If improvement in milk volume is the measurable response, a breakeven point can be calculated. For example, an additive that raises feed costs 10 cents per day is used. If milk is valued at 12 cents per pound, every cow must produce 0.84 pounds more milk to cover the extra cost associated with the additive. Another consideration is if all the cows receive the additive, but only cows fresh for less than 100 days respond. These responding cows must cover the costs for all cows (responsive and nonresponsive). One guideline is that an additive should return \$2 or more for each dollar invested to cover nonresponsive cows and field conditions that could minimize the anticipated response.

Remember, it is difficult to assess management practices that acutely alter milk production. Research is essential to determine if experimentally measured responses can be expected in the field. Rely on research studies conducted under controlled and unbiased conditions that use an experimental protocol similar to field conditions and that have statistically analyzed results.

Results obtained on individual farms are the economic payoff. Managers and consultants must use a database to accurately compare and measure responses. Several tools to measure results include DHIA milk production records, reproduction summaries, somatic cell count data, dry matter intake, heifer growth charts, body condition scores and herd health profiles. These tools will enable managers and consultants to critically evaluate the effectiveness of selected additives.

Heart of America Dairy Herd Improvement Summary

	Quartiles				Your Herd
	1	2	3	4	
Ayrshire					
Rolling Herd Average	17,025	15,295	13,665	11,370	
Summit Milk Yield 1st	60.0	57.5	48.0	21.5	
Summit Milk Yield 2nd	37.5	62.5	59.0	26.0	
Summit Milk Yield 3rd	41.5	71.5	66.5	60.0	
Summit Milk Yield Avg.	64.0	65.0	57.5	58.5	
Income/Feed Cost	935.0	782.0	715.5	674.5	
SCC Average	263.0	142.0	445.5	128.5	
Days to 1st Service	61.5	65.5	84.0	68.0	
Days Open	176.5	124.5	171.5	84.5	
Projected Calving Interval	15.0	13.3	14.8	7.40	
Brown Swiss					
Rolling Herd Average	19,666	16,729	15,446	13,840	
Summit Milk Yield 1st	63.8	54.8	50.5	50.1	
Summit Milk Yield 2nd	64.3	64.4	63.5	60.8	
Summit Milk Yield 3rd	83.3	64.2	68.6	65.0	
Summit Milk Yield Avg.	75.1	64.5	61.3	59.5	
Income/Feed Cost	1,858	1,100	1,004	875	
SCC Average	445	354	433.0	362.8	
Days to 1st Service	102	89.7	77.1	86.0	
Days Open	191.6	182.4	168.8	202.0	
Projected Calving Interval	15.5	15.2	14.7	15.8	
Guernsey					
Rolling Herd Average	15,556	14,086	13,535	12,271	
Summit Milk Yield 1st	56.0	43.0	50.0	43.0	
Summit Milk Yield 2nd	69.0	65.0	61.0	58.3	
Summit Milk Yield 3rd	68.5	64.5	67.0	57.0	
Summit Milk Yield Avg.	63.5	60.5	57.5	53.6	
Income/Feed Cost	1,070	1,330	678.5	809.6	
SCC Average	524.5	156	468.5	247	
Days to 1st Service	102.0	42.5	41.0	107.3	
Days Open	165.5	240.5	220	217.3	
Projected Calving Interval	14.6	17.1	16.4	16.3	
Holstein					
Rolling Herd Average	23,306	20,226	18,053	14,711	
Summit Milk Yield 1st	73.5	66.4	60.9	52.0	
Summit Milk Yield 2nd	93.4	83.5	75.1	62.6	
Summit Milk Yield 3rd	98.4	88.8	80.7	67.9	
Summit Milk Yield Avg.	87.3	79.3	72.7	62.4	
Income/Feed Cost	1,618	1,399	1,149	868	
SCC Average	379.3	404.9	446.3	554.9	
Days to 1st Service	93.2	93.2	97.6	90.5	
Days Open	164	167.6	181.6	202.1	
Projected Calving Interval	14.6	14.7	15.1	15.8	
Jersey					
Rolling Herd Average	17,135	14,671	13,632	11,801	
Summit Milk Yield 1st	53.2	47.4	43.7	43.4	
Summit Milk Yield 2nd	65.6	62.8	53.0	51.1	
Summit Milk Yield 3rd	69.0	63.0	52.8	51.3	
Summit Milk Yield Avg.	63.0	57.3	53.8	49.0	
Income/Feed Cost	1,560	1,316	1,033	828	
SCC Average	256.7	349.9	551.7	454.2	
Days to 1st Service	83.5	94.9	71.8	100	
Days Open	136	152.8	157.8	148.9	
Projected Calving Interval	13.7	14.2	14.3	14.1	
Milking Shorthorn					
Rolling Herd Average	14,508	13,586	14,138	9,517	
Summit Milk Yield 1st	52.0	51.0	45.0	57.0	
Summit Milk Yield 2nd	73.0	60.0	58.0	—	
Summit Milk Yield 3rd	69.0	67.0	65.0	63.0	
Summit Milk Yield Avg.	60.0	60.0	59.0	62.0	
Income/Feed Cost	852	961	1030	—	
SCC Average	230	205	373	154	
Days to 1st Service	67.0	91.0	81.0	—	
Days Open	155	131	177	168	
Projected Calving Interval	14.3	13.5	13.7	14.8	

Hay Prices*—Kansas

	Location	Quality	Price (\$/ton)
Alfalfa	Southwestern Kansas	Supreme	120-130
Alfalfa	Southwestern Kansas	Premium	120-130
Alfalfa	Southwestern Kansas	Good	—
Alfalfa	South Central Kansas	Supreme	100-110
Alfalfa	South Central Kansas	Premium	100
Alfalfa	South Central Kansas	Good	—
Alfalfa	Southeastern Kansas	Supreme	—
Alfalfa	Southeastern Kansas	Premium	100
Alfalfa	Southeastern Kansas	Good	—
Alfalfa	Northwestern Kansas	Supreme	100-105
Alfalfa	Northwestern Kansas	Premium	100
Alfalfa	Northwestern Kansas	Good	—
Alfalfa	North Central/East Kansas	Supreme	110-120
Alfalfa	North Central/East Kansas	Premium	100-110
Alfalfa	North Central/East Kansas	Good	—

Supreme = over 180 RFV (less than 27 ADF)
 Premium = 150–180 RFV (27–30 ADF)
 Good = 125–150 RFV (30–32 ADF)

Source: USDA Kansas Hay Market Report, April 17, 2001.

Hay Prices—Oklahoma


	Location	Quality	Price (\$/ton)
Alfalfa	Central/Western, OK	Premium	110-130
Alfalfa	Central/Western, OK	Good	100-115
Alfalfa	Panhandle, OK	Premium	105-130
Alfalfa	Panhandle, OK	Good	100-120

Source: Oklahoma Department of Agriculture, April 12, 2001

Feed Stuffs Prices

	Location	Price (\$/ton)
Blood Meal	Central US	391
Canola Meal		140-145
Corn Gluten Feed	Kansas City	55-60
Corn Gluten Meal	Kansas City	235-240
Corn Hominy	Kansas City	48-52
Cotton Seed Meal	Kansas City	160-165
Whole Cotton Seed	Memphis	105
Distillers Grains	Central Illinois	98-110
Pork—Meat and Bone Meal	Texas Panhandle	150-160
SBM 48%	Kansas City	158-164
Sunflower Meal		90
Wheat Middlings	Kansas City	47-50

Source: USDA Feedstuff Market Review, April 11, 2001



COOPERATIVE EXTENSION SERVICE
U.S. DEPARTMENT OF AGRICULTURE
KANSAS STATE UNIVERSITY
MANHATTAN, KANSAS 66506

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE. \$300

Dairy Lines is jointly published for dairy producers by the Department of Animal Sciences and Industry, K-State Research and Extension, and the Department of Animal Science, Oklahoma Cooperative Extension Service. For more information or questions, please contact 785.532.5654 (K-State) or 405.744.6058 (OSU).

Kansas State University
K-State Research & Extension
Department of Animal Sciences and Industry
Call Hall, Room 139
Kansas State University
Manhattan, Kansas 66506

KSU, County Extension Councils and U.S. Department of Agriculture Cooperating.
All educational programs and materials available without discrimination on the basis of color, race, religion, national origin, sex, age, or disability.

The Department of Animal Sciences and Industry at Kansas State University greatly appreciates the sponsor(s) of the Dairy Lines Newsletter. These sponsorships in no way imply the Department's endorsement of the products and services offered by the sponsors. The Department welcomes inquiries from other individuals, associations and firms that may be interested in cosponsoring this publication.




John Smith
Extension Specialist
Dairy Science
K-State



Mike Brouk
Extension Specialist
Dairy Science
K-State



Dan Waldner
Extension Specialist
Dairy Science
Oklahoma State



Dairy Lines



DAIRY RESEARCH AND EXTENSION NEWS
K-State Research and Extension
and Oklahoma State University