

For additional synchronization protocols visit BeefRepro.org
 Use of sexed semen can result in conception rates that are decreased compared to conventional semen used in the same situation

HEAT DETECTION


Sexed semen can be used on any female observed in estrus (heat) and following synchronization with any protocol on the Cow or Heifer Protocol Sheet.
 For best results with sexed semen perform AI 16 to 22 hr after detecting female in estrus.
 Use of estrus detection aids (EDA) is highly recommended.

FIXED-TIME AI

Time for Fixed-time AI (TAI) should be considered as the approximate average time of insemination based on the number of females to inseminate, labor, and facilities.
 To optimize pregnancy success, it is recommended to use sexed semen on females that have exhibited estrus before TAI and conventional semen on females that have not exhibited estrus.


HEIFERS ONLY 7-day CO-Synch+CIDR®

Perform TAI with GnRH at 72 ± 2 hr after PG.




COWS 7&7 Synch

Perform TAI with GnRH at 66 ± 2 hr after CIDR removal. *All cows can also be inseminated with either sexed or conventional semen.*



HEIFERS ONLY PG-7 7-day CO-Synch+CIDR®


Perform TAI with GnRH 72 ± 2 hr after second PG. *All heifers can also be inseminated with either sexed or conventional semen.*




SPLIT-TIME AI (STAI)

Perform first TAI 66 ± 2 hr after PG in estrous females. GnRH administered at second TAI only in non-estrous females. Females detected in estrus by first or second TAI receive sexed semen.
 Females not in estrus receive conventional semen at second TAI.

HEIFERS ONLY 14-day CIDR®-PG & STAI



COWS & HEIFERS 7-day CO-Synch+CIDR® & STAI



EDA

Aervoe™, AIPaint™, Detect-her™, Estroject™, Kamar™, Mark-her™, Paintstik™, Quick Shot™, Tell Tail™, Twist-Stik™

GnRH

Cystorelin®, Factrel®, Fertagyl®, OvaCyst®, GONABreed®

PG

estroPLAN®, Estrumate®, Lutalyse®, Lutalyse® HighCon, ProstaMate®, SYNCHSURE™



SCAN