

## 260 Effects of iron administration timing on pre-weaning performance and hematological criteria in pigs.

Hayden E. Williams<sup>1</sup>, Ryan T. Maurer<sup>1</sup>, Brittany Carrender<sup>2</sup>, Joel M. DeRouchey<sup>1</sup>, Jason C. Woodworth<sup>3</sup>, Steve S. Dritz<sup>1</sup>, Mike D. Tokach<sup>1</sup>, Robert D. Goodband<sup>1</sup>, Kyle Coble<sup>2</sup>, Andrew Holtcamp<sup>4</sup>, <sup>1</sup>*Kansas State University*, <sup>2</sup>*JBS Live Pork*, <sup>3</sup>*Department of Animal Sciences & Industry, College of Agriculture, Manhattan, KS 66506*, <sup>4</sup>*CEVA Animal Health*

Newborn pigs (n=1,892; 1.5 kg BW) were used in a 20-d study evaluating the effects of Fe injection timing after birth on preweaned pig performance and blood criteria. A total of 172 litters were used. One d after farrowing, piglets were weighed, and 11 pigs within each litter were allotted to 1 of 6 treatments in a CRD. Treatments consisted of pigs receiving no Fe injection or 200-mg of injectable Fe (GleptoForte, Ceva Animal Health, Lenexa, KS) provided on d 1, 3, 5, or 7 of age, or 200-mg on d 1 plus 200-mg on d 12. 1 pig/litter received no Fe injection and 2 pigs/litter were placed on all other treatments. Piglets were weighed on d 1 and 20 after birth to determine growth performance and bled on d 20 to determine Fe status. Increasing the age that piglets received the Fe injection tended to decrease (linear;  $P=0.080$ ) ADG. Not providing an Fe injection decreased ( $P=0.0001$ ) overall ADG and d 20 BW compared to all other treatments. Hemoglobin and Hct decreased (linear;  $P < 0.05$ ) with increasing age when pigs received an Fe injection. There was no evidence of differences ( $P > 0.10$ ) between the pigs receiving a 200-mg injection on d 1 and d 12 compared to those receiving the Fe on d 1 only. Pigs not provided an Fe injection had decreased ( $P=0.0001$ ) Hb and Hct values compared to pigs receiving an Fe injection. Pigs receiving the 200-mg injection on d 1 and 12 had increased ( $P=0.0001$ ) Hb and Hct values compared to pigs receiving 200-mg on d 1 only. Results suggest that providing a 200-mg Fe injection within 7 d after farrowing is sufficient for optimizing preweaning growth performance. The additional 200-mg Fe injection at d 12 did not influence growth performance but does increase Hb and Hct at weaning.

**Table 1. Day of iron administration effects on pre-weaning performance and hematological criteria in pigs**

Item	None <sup>2</sup>	Fe injection day <sup>1</sup>				
		1	3	5	7	1 and 12
BW, kg <sup>3</sup>						
d 20	5.1	6.1	6.0	6.1	6.0	6.1
SEM	0.08	0.06	0.06	0.06	0.06	0.06
ADG, g <sup>3,4</sup>						
d 1 to 20	192	248	244	247	240	249
SEM	3.73	2.76	2.78	2.79	2.81	2.80
Hb, g/dL <sup>3,5</sup>						
d 20	5.0	11.1	11.3	10.7	10.3	12.2
SEM	0.32	0.20	0.20	0.20	0.20	0.20
Hct, % <sup>3,5</sup>						
d 20	15.2	32.6	33.3	31.7	30.4	36.1
SEM	1.06	0.64	0.63	0.64	0.65	0.64

<sup>1</sup>200 mg of Fe (GleptoForte, Ceva Animal Health, LLC., Lenexa, KS) administered on d 1, 3, 5, 7, or d 1 and 12 after farrowing.

<sup>2</sup>Negative control with pigs receiving no iron injection.

<sup>3</sup>0 vs. others ( $P = 0.0001$ ).

<sup>4</sup>Day main effect (linear;  $P = 0.080$ ).

<sup>5</sup>Day main effect (linear;  $P < 0.05$ ).

**Keywords:** iron, performance, preweaning